



# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING

# TRUMP<sup>†</sup>

## SPRAY OIL

**ACTIVE CONSTITUENT:**  
**830 g / L PARAFFINIC OIL**

UNSULPHONATED RESIDUE NOT LESS THAN 98.0%

**A NARROW RANGE EMULSIFIABLE PARAFFINIC OIL  
FOR THE MANAGEMENT OF PESTS AND AS AN ADJUVANT  
OR ULV CARRIER FOR USE WITH PESTICIDES AND  
DEFOLIANTS AS PER DIRECTIONS FOR USE**

**IMPORTANT:** Read the attached leaflet before use.

**VICCHEM**  
*The Right Chemistry*



ACO CERT. NO. 11155

PEEL  
HERE

# TRUMP<sup>†</sup>

## SPRAY OIL

### STORAGE AND DISPOSAL

Store in the closed, original container, in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsate to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in an approved waste facility. If an approved waste facility is not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. DO NOT burn empty containers or product. Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

### SAFETY DIRECTIONS

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles.

### First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

### SAFETY DATA SHEET

Additional information is listed in the Safety Data (MDS).

### Hazard Statements

Harmful if swallowed. Causes serious eye irritation

### Precautionary Statements

Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. If swallowed rinse mouth. Call a POISON CENTER or doctor if you feel unwell. Wear eye protection. IF IN EYES: Rinse cautiously with water for several minutes. Re-move contact lenses if present and easy to do. Continue rinsing. If eye irritation persists get medical advice

### APVMA Approval No.

59092/145870



**Victorian Chemical  
Company Pty. Limited**  
83 Maffra Street, Coolaroo,  
Victoria 3048, Australia  
Telephone: (03) 9301 7000  
Facsimile: (03) 9309 7966  
Email: products@vicchem.com  
Website: www.vicchem.com



**NOTICE:** Victorian Chemical Company Pty. Limited accepts no responsibility for any consequences whatsoever arising from the use of this product except responsibility imposed under any applicable law. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this product on these conditions.

† Trademark used under licence. \* Third Party Trademark.



For applicable packs

919121B-8

**Restraints:**

DO NOT spray when temperatures exceed 32°C, (except where noted for citrus) or when soil is dry and trees are suffering from moisture stress.

**SPRAY DRIFT RESTRAINTS**

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment

designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application.

Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise

**DIRECTIONS FOR USE:**

Fruits, vegetable and berries

Crop	Pest	State	Application Rate	Critical Comments
<b>Almonds, Peaches, Nectarines</b>	Aphids (Eggs), Mites (Eggs)	NSW, ACT, Vic, SA, WA, Tas only	3L/100L	Dormant or delayed dormant
	San Jose Scale	NSW, ACT, Vic, SA, WA only	2L/100L	Dormant
	Mites and Scales	NSW, ACT, Vic, SA, WA only	1L to 2L/100L	Summer (Foliar or Cover) or Post Harvest.
<b>Apples</b>	Mites (Eggs), including European Red Mite, Scales (Hard & Soft)	NSW, ACT, Vic, SA, WA, Tas	2L/100L	Dormant or delayed dormant to half green tip
		Qld	3L/100L	
	Mites	NSW, ACT, Vic, SA, WA, Tas	1L/100L	Summer (Foliar or Cover) or Post Harvest
<b>Apricots, Cherries</b>	San Jose Scale	NSW, ACT, Vic, SA, WA only	2L/100L	Dormant
		Qld	3L/100L	
	Mites and Scales	NSW, ACT, Vic, SA, WA, Tas only	1L to 1.5L/100L	Summer (Foliar or Cover) or Post Harvest. Applications should not be made over oil sensitive varieties. Do not apply to trees lacking adequate moisture.
<b>Asparagus, Beans, Beet, Corn, Cucurbits, Peppers/ Capsicum, Radish, Squash, Tomatoes</b>	Aphids Mites Thrips Leafhopper	All States (Except Qld for beans, tomato, corn and peppers / capsicums)	1 to 2L/100L	As needed.

Crop	Pest	State	Application Rate	Critical Comments
<b>Avocado</b>	Scale Insects	Qld	1.2L to 1.6L/100L	Apply when heavy scale populations occur on stem, foliage or fruit. Do not apply if trees need watering. Application is most effective against young crawler stages.
		NSW, ACT, WA	2L/100L	
<b>Bananas</b>	Cercospora Leaf Spot	NSW, ACT	4L/ha	Apply in combination with Mancozeb at 2.2-4.5 kg/ha in 200 to 500L of water. Apply at 3-4 week intervals, from December until late April. In North Queensland the interval may be reduced to 10-14 days in the wet season.
		Qld	3.5L/ha	
	Cercospora Leaf Spot, Leaf Speckle	NSW, ACT, Sth Qld, WA	5L/ha	Apply in combination with Tilt* 250 EC. Commence spraying in December and apply 4 to 5 times at 21 day intervals. Later use alternative fungicides if needed. Spray both sides of leaves.
	Cercospora Leaf Spot, Leaf Speckle, Cordana Leaf Spot	Nth Qld	5L/ha	Apply in combination with Tilt* 250 EC. Spray every two weeks from the start of the wet season in December and make a maximum of 6 applications. Later use alternative fungicides if needed. Spray both sides of leaves.
	Scale Insects	Qld, NSW, ACT, WA	2L/100L	Apply when heavy scale populations occur on stem, foliage or fruit. Do not apply if trees need watering. Application is most effective against young crawler stages.
<b>Blueberries</b>	Mites, Scales	All States	2L/100L	Foliage.
			3L/100L	Dormant.
<b>Corn (Sweet and Field) Sugar Beet</b>	Aphids, Mites, Whitefly Bugs (Immature)	NSW, ACT, SA, WA, Tas only	2L/100L	As needed
<b>Citrus</b>	Redscale	Vic, SA, WA	1 to 2L/100L	Apply in December and/or February - March
		NSW, ACT	1.3 to 2L/100L	
		Qld	1L/100L	Apply late November/early December and late January/early February if required – apply in combination with Methidathion 50mL/100L.
	White Wax Scale, Pink Wax Scale	Vic, WA	1 to 2L/100L	Apply in December / early January Period
		NSW, ACT	2L/100L	
		Qld	1L/100L	Apply late November/early December and late January/early February if required. Apply in combination with methidathion 50mL/100L, carbaryl 70g/100L, promecarb 50g/100L, soda ash 500-750g/100L or washing soda 500-750g/100L.
	White Wax Scale, Pink Wax Scale, Black Scale	Tas, WA	1.8 to 2.5L/100L	Apply mid December/early January. Repeat in February.
	Soft Brown Scale	Tas, WA	1.8 to 2.5L/100L	Apply in January / April.

Crop	Pest	State	Application Rate	Critical Comments
Citrus	Circular Black Scale	QLD	500mL to 600mL/100L	Apply in late Nov/early Dec and in late Jan/early Feb if required
		WA	500mL to 1L/100L	Apply in late Nov/early Dec and in late Jan/early Feb if required
	Citrus Leaf-miner	All States	250mL to 1L/100L (for Citrus Leaf-miner Control only) 500mL to 1L/100L (when combining with Scale & Black Spot Control)	Apply thoroughly and evenly to flush growth, wetting both sides of all immature leaves to point of run-off. Spray every 5 to 14 days during flush periods, commencing as soon as pest's silvery mites are observed on young leaves. The interval will be shorter during summer and early autumn. Cease spraying when leaves become mature. Four sprays in one year should be sufficient to achieve control and no more than 8 sprays should be applied.
	Broad Mites	All states	250mL/100L + 15 or 25 mL Vetimec* Miticide/Insecticide or similar	Apply in accordance with the Vetimec* Miticide/Insecticide or similar label.
	Brown Citrus rust mites	All states	700mL to 1L/ha	Apply Nov/Dec and again in Feb/Mar if required.
	Citrus Red Mites	NSW Only	700mL to 1L/ha	Apply Aug/Sep and again in Mar/Apr if required.
	Citrus Rust Mite	All States	250mL/100L + 15 or 25 mL Vetimec* Miticide/Insecticide or similar	Apply in accordance with the Vetimec* Miticide/Insecticide or similar label.
		QLD Only	700mL to 1L/ha	Apply Nov/Dec and again in Feb/Mar if required.
		NSW, Vic, SA, WA Only	700mL to 1L/ha	Apply in Feb/Mar.
	Black Spot	All States	600mL/100L + copper oxychloride	Apply at half to 1/4 petal fall. Consult copper fungicide label for suitable local rate and intervals for repeat application. Ensure thorough coverage is achieved throughout the canopy.
		QLD Only	600mL/100L + 200g Dithane* Rainshield Fungicide or Penncozeb* 750DF Fungicide or similar.	Apply in accordance with registered Dithane* Rainshield Fungicide or Penncozeb* 750DF Fungicide or similar.
		NSW, Tas, Vic & SA only	250mL/100L + 200g Dithane* Rainshield Fungicide or Penncozeb* 750DF Fungicide or similar.	Apply in accordance with registered Dithane* Rainshield Fungicide or Penncozeb* 750DF Fungicide label or similar.
	Melanose, Smoky Blotch, Scab	All States	600mL/100L + copper oxychloride	Apply in accordance with registered copper oxychloride label. Ensure thorough coverage is achieved throughout canopy.

Crop	Pest	State	Application Rate	Critical Comments
Citrus	Brown Spot in Mandarins	QLD Only	600mL/100L + 375g copper oxychloride + 500g zinc sulphate heptahydrate + 140g caustic soda	Apply in accordance with registered copper oxychloride label.
			625mL/100L + 200 to 300g copper oxide	Apply in accordance with registered copper oxide label.
	Brown Rot	QLD Only	625mL/100L + 200 to 300g copper oxide	Apply in accordance with registered copper oxide label.
Custard Apple	Scale Insects	Qld	1 to 1.7L /100L	Apply when heavy scale populations occur on stem, foliage or fruit. Do not apply if trees need watering. Application is most effective against young crawler stages
		NSW, ACT, WA	2L/100L	
Grapes	Mealybugs	NSW, ACT, Qld, SA, WA, Tas only	4 to 6L /100L with 306L to 460L Water per Hectare	Dormant - A tank mix using insecticides improves control. Air carrier or Air blast type sprayers strongly recommended to avoid plant injury and reduce effect on bloom of table grapes
	Grapevine Scale	NSW, Vic, SA, WA, Qld only	3L/100L	Apply as a post pruning spray in mid-winter when vines are fully dormant.
		Tas only	2L/100L	
	Powdery Mildew	All States	1L/100L	Commence spraying just after woolly bud stage. Follow-up sprays should be applied every 14-21 days till bunch closure.
Kiwi Fruit	Scale Insects	Qld	1 to 1.6L/100L	Apply when heavy scale populations occur on stem, foliage or fruit. Do not apply if trees need watering. Application is most effective against young crawler stages.
		NSW, ACT, WA	2L/100L	
Mango	Scale Insects	Qld, NSW WA, NT only	1 to 2L/100L	Apply after fruit picking when heavy scale populations occur on stem and foliage.
	Wax Scale			
	Mango Plant Hopper			
Olives	Scales (Black Scale, Soft Brown, etc)	All states	1L/100L	Best results are achieved when applied to crawlers in spring. Minimum of 1000L water per Ha
	Black Scale, Olive Scale, California Red Scale	All states	1.8 to 2.5L/100L	DO NOT spray in winter as product will not rapidly break down. To prevent fruit spotting in green table olives, DO NOT spray for 6 weeks prior to harvest.

Crop	Pest	State	Application Rate	Critical Comments
<b>Passion Fruit</b>	Red Scale, Hemispherical Scale Suppression of Passionvine Mealy Bug	All States	500mL - 1L/100L	Please refer to the General Instructions under “Crop Safety — Passionfruit” for full details on crop safety. Apply via a low volume rate of 1000L per hectare. A minimum interval of 14 days should elapse between sprays in the first month with additional applications applied at monthly intervals for up to 6 months. Good coverage is essential for Red Scale.
<b>Pears</b>	European Red Mite, Eggs, Scales	NSW, ACT, Vic, SA, WA	1 to 2L /100L	Dormant or delayed dormant (up to and including petal fall)
		Tas only	2L/100L	
		Qld	3L/100L	
	Two Spotted Spider Mite	NSW, ACT, Vic, SA, WA, Tas only	1L/100L	Summer (Foliar or Cover) or Post Harvest.
	Brown Mite, Pear Rust Mite	NSW, ACT, Vic, SA, WA, Tas only	1 to 2L/100L	
	Pear Leaf Blister Mite	NSW, ACT, Vic, SA, WA, Tas only	1.5 to 2L/100L	
<b>Pecans</b>	Aphids (Eggs), Scales, Mites (Eggs)	NSW, ACT, Qld, SA, WA, Tas only	3L/100L	Dormant
	Aphids, Mites	NSW, ACT, Qld, SA, WA, Tas only	1 to 1.5L/100L	Summer (Foliar or Cover) or Post Harvest.
<b>Plums Prunes</b>	Aphids (Eggs), Scales, Mites (Eggs)	NSW, ACT, SA, WA, Tas only	1.5L to 2L/100L	Dormant or delayed dormant. To half green tip.
	Mites, Scales	NSW, ACT, Vic, SA, WA, Tas only	1 to 1.5L/100L	Summer (Foliar or Cover) or Post Harvest. Do not apply over fresh market fruits after bloom starts to form as the oil will remove the bloom. Application should not be made over sensitive varieties.
<b>Strawberries</b>	Aphids, Mites	NSW, ACT, SA, WA, Tas only	1L/100L	Dormant and Foliage.

#### Ornamental, Flowers, Foliage, Trees and Shrubs

Crop	Pest	State	Application Rate	Critical Comments
<b>Shade Trees* and Shrubs including Conifers, Deciduous Broadleaf Evergreens and Woody Ornamentals, Ornamentals Trees*, Shrubs along city streets, other rights of way including Deciduous and Broadleaf Evergreens.</b>	Aphids, Scales (Immature) Spider Mites, Whitefly (Immature)	NSW, ACT, Vic, SA, WA, Tas only	2 to 4L/100L (Winter)	Winter Dormant period as needed for plants that become dormant in Winter.
			1 to 3L/100L (Summer)	Summer (Foliar or Cover) as needed.

Crop	Pest	State	Application Rate	Critical Comments
<b>Flowers and Foliage Plant including Roses and other Flowering Shrubs, Foliage Ornamentals and Bedding Plants</b>	Aphids, Scales (Immature) Spider Mites, Whitefly (Immature)	NSW, ACT, Vic, SA, WA, Tas only	2 to 3L/100L (Winter)  1 to 2L/100L (Summer)	Winter Dormant period as needed for plants that become dormant in Winter.  Summer (Foliar or Cover) as needed.
<b>Chrysanthemum Diffenbachia Dracaena Ferns Gardenias Jade Plant Palms Philodendron</b>	Aphids Scales Whitefly	All States	10 to 20mL/L OR 1 to 2L/100L	Do not apply to plants in direct sunlight behind glass. Do not use on Coconut Palms and Maidenhair Ferns. Chrysanthemum blooms have shown phytotoxic symptoms at the higher rate. Applicator should conduct a test for phytotoxicity by treating a few specimens before making a large-scale application. Protect floor, floor coverings and furnishings from overspray.
<b>Azalea Begonia Camellia Chrysanthemum Crown of Thorns</b>	Aphids Mealybugs Scale (Soft & Hard) Thrips	All States	10 to 20mL/L OR 1 to 2L/100L	Do not apply to plants in direct sunlight behind glass.
<b>Diffenbachia*</b>	Whitefly (Immature)		10mL/L OR 1L/100L	
<b>Easter Lily Fern Gardenia Hibiscus Foliage Jade Plant Palm Philodendron</b>	Reduces number of Spider Mites		10 to 20mL/L OR 1 to 2L/100L	
<b>Poinsettia</b>			10mL/L OR 1L/100L	
<b>Reiger Begonia Zinnias</b>			10 to 20mL/L OR 1 to 2L/100L	
<b>Hardy Plants</b>	Leaf Polish	All States	10mL/L OR 1L/100L	
<b>Christmas Trees</b>	Aphids, Scales (Soft & Hard) (Immature) Spider Mites	NSW, ACT, Vic, SA, WA, Tas only	2 to 4L/100L (Winter)  1 to 3L/100L (Summer)	Winter  Summer
<b>CAUTION:</b> Spray no more than 4 times during growing season with two weeks minimum application interval. Do not spray when buds are fully opened and shoot elongation is occurring. Do not spray when there is no obvious moisture in the leaves or the plant is under stress. Avoid spraying open blooms. Bleaching and spotting has been observed with open blooms of certain plants. Oil removes the glaucous (blue) bloom from some evergreens. Always use lower dosage or test spray oil sensitive plants first – do not spray walnut foliage.				



**Field Crops**

Crop	Pest	State	Application Rate	Critical Comments
Cotton	Defoliation Aid	NSW, ACT, Qld, WA	500mL to 1L/100 L	Apply in combination with thidiazuron (eg DROPP*) or thidiazuron + diuron (eg. DROPP* Ultra) defoliation products. The high rate may help late-season defoliation or when defoliation conditions are not ideal.
	Carrier for certain ULV insecticides	NSW, Qld only	3 to 5L/ha	Apply in combination with ULV insecticide as specified on the insecticide label to ensure insecticide reaches its target.

**For use as an adjuvant with herbicides, fungicides and insecticides**

Crop	Pest	State	Application Rate	Critical Comments
Wetting Agent if recommended on Herbicide label	Weeds as specified on herbicide label	All States	250 to 500mL/100L	TRUMP Spray Oil may be used as a crop oil in a spray mixture with selective herbicides to improve their wetting and penetrating properties.
			2L/ha	
Wetting agent/ adjuvant /carrier as recommended on fungicide and insecticide labels	Refer to particular pesticide label	All States	500mL to 5L/100L	TRUMP Spray Oil acts as a carrier /adjuvant /wetting agent which may improve the performance of certain pesticides. Always check pesticide label for correct rates.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**Withholding Periods:**

NOT REQUIRED when used as directed. When used in tank mix or as an adjuvant, adhere to the withholding period specified on the label of the companion product(s) used

**GENERAL INSTRUCTIONS**

TRUMP Spray Oil controls Aphids, Lace Bugs, Leaf Hoppers, Mealybugs, Mites, Plant Bugs, Scale, Whitefly, Eggs of Aphids and Mites on vegetables, Fruit Trees, Nut Trees, certain field crops, Shrubs, Plantation Trees, Greenhouse Plants, Ornamental Foliage Plants and Flowers. TRUMP Spray Oil can be used in combination with certain fungicides, herbicides and insecticides as a wetting agent/adjuvant/carrier as specified on the products label.

**MIXING INSTRUCTIONS**

1. Add water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides.
  - a. If wettable powder formulation – mix water and powder thoroughly so that powder is totally suspended in the water before the oil is added.
  - b. If other pesticide to be added is an emulsifiable formulation, do so after the oil and water has been thoroughly mixed.
3. Add oil under agitation when tank is three quarters full. Top off with water to form a milky solution.
4. Maintain agitation until solution is completely used.
5. In a small equipment lacking agitators, stir or shake diluted spray frequently during applications.
6. It is important for users to read and follow all instructions on the labels of the proposed tank mixed products.
7. Flush fluid in sprayer hose lines back into tank reservoir if fluid is allowed to stand for more than 20 minutes.

## **APPLICATION INSTRUCTIONS**

The target must be completely covered in spray.

Oil residue on the plants surface often acts as a feeding and oviposition deterrent.

However, the primary target is the pest itself as TRUMP Spray Oil is a contact pesticide.

Dilute applications. (Greater than 1400L spray per hectare in horticultural crops) in most cases ensure best coverage.

Speed of travel is extremely important, tractor speed from 1.6 kph to 5.4 kph is recommended, depending on crop size and target pest.

Do not spray when shade temperatures are or exceed 32°C.

## **COMPATIBILITY**

TRUMP Spray Oil can be used with such pesticides as NPV's, Bt's, Pyrethroids, Abamectin, Tracer II\*, Mancozeb and Folicur\* as stated on their respective directions for use labels.

TRUMP is compatible with certain grain seed treatment chemicals.

Do not use spray oils with Dimethoate.

If possible, either keep the spray equipment used for these compounds separate from the equipment used for TRUMP Spray Oil, or make sure that the sprayer is thoroughly cleaned, so that no residue from these compounds remain.

Do not use with Carbaryl on deciduous fruit trees.

## **TANK MIXTURES**

TRUMP Spray Oil may be mixed with other pesticides to improve the level of kill or enhance coverage. User should read and follow all instructions on the label of the proposed tank mixed product.

## **TIMING THE TREATMENT**

Applicator must determine the precise timing to fit local growth and climate conditions. Do not exceed maximum rates or apply more often than recommended.

May be used up to one day before harvest.

## **CROP SAFETY — PASSIONFRUIT**

The sensitivity of this product has not been tested on all Passionfruit varieties, therefore it is recommended that a small number of plants be treated in the first instance before treating the whole crop.

DO NOT exceed an application rate of 0.5% or 4 applications per year for oil sensitive varieties such as those from the Panama line (Passionfruit only).

## **Precautions:**

Do not allow entry into treated area until spray is dried. When using together with other products, consult their re-entry period. The longer re-entry period should be followed

## **Protections:**

### **PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**

Dangerous to fish. Do not contaminate streams, rivers or watercourses with the chemical or used containers.

### **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Do not apply pesticides or other chemicals incompatible with oil, within 4 weeks before or after application of any other spray oils, as crop damage may occur. Do not apply within 3 weeks before or after a spray containing sulphur.

## **Storage and Disposal:**

Store in the closed, original container, in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsate to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler

or designated collection point. If not recycling, break, crush or puncture and bury empty containers in an approved waste facility. If an approved waste facility is not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. DO NOT burn empty containers or product. Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage

**Safety Directions:**

Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray wear elbow-length PVC gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles

**First Aid Instructions:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

**SAFETY DATA SHEET**

Additional information is listed in the Safety Data (MDS).

**Hazard Statements**

Harmful if swallowed. Causes serious eye irritation

**Precautionary Statements**

Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. If swallowed rinse mouth. Call a POISON CENTER or doctor

if you feel unwell. Wear eye protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do.

Continue rinsing. If eye irritation persists get medical advice

**APVMA** Approval No. 59092/145870

\* Third Party Trademark

† Trademark used under licence

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# TRUMP<sup>†</sup>

## SPRAY OIL

**ACTIVE CONSTITUENT : 830 g / L PARAFFINIC OIL**  
**UNSULPHONATED RESIDUE NOT LESS THAN 98.0%**

A NARROW RANGE EMULSIFIABLE PARAFFINIC OIL FOR  
THE MANAGEMENT OF PESTS AND AS AN ADJUVANT OR  
ULV CARRIER FOR USE WITH PESTICIDES AND  
DEFOLIANTS AS PER THE DIRECTIONS FOR USE.

**IMPORTANT: Read the attached leaflet before use.**



**Victorian Chemical  
Company Pty. Limited**  
83 Maffra Street, Coolaroo,  
Victoria 3048, Australia  
Telephone: (03) 9301 7000  
Facsimile: (03) 9309 7966  
Email: [products@vicchem.com](mailto:products@vicchem.com)  
Website: [www.vicchem.com](http://www.vicchem.com)



**NOTICE:** Victorian Chemical Company Pty.  
Limited accepts no responsibility for any  
consequences whatsoever arising from the use  
of this product except responsibility imposed  
under any applicable law. The buyer assumes all  
risk and liability resulting from improper  
handling, storage or use and accepts and uses  
this product on these conditions.

\* Third party trademark



For applicable packs

919121B-8